



**Pharming Group N.V.**


24<sup>th</sup> Annual Needham  
Virtual Healthcare Conference

**April 7-10, 2025**

NASDAQ: **PHAR** | EURONEXT Amsterdam: **PHARM**

*This presentation may contain forward-looking statements. Forward-looking statements are statements of future expectations that are based on management's current expectations and assumptions and involve known and unknown risks and uncertainties that could cause actual results, performance, or events to differ materially from those expressed or implied in these statements. These forward-looking statements are identified by their use of terms and phrases such as "aim", "ambition", "anticipate", "believe", "could", "estimate", "expect", "goals", "intend", "may", "milestones", "objectives", "outlook", "plan", "probably", "project", "risks", "schedule", "seek", "should", "target", "will" and similar terms and phrases. Examples of forward-looking statements may include statements with respect to timing and progress of Pharming's preclinical studies and clinical trials of its product candidates, Pharming's clinical and commercial prospects, and Pharming's expectations regarding its projected working capital requirements and cash resources, which statements are subject to a number of risks, uncertainties and assumptions, including, but not limited to the scope, progress and expansion of Pharming's clinical trials and ramifications for the cost thereof; and clinical, scientific, regulatory, commercial, competitive and technical developments. In light of these risks and uncertainties, and other risks and uncertainties that are described in Pharming's 2024 Annual Report and the Annual Report on Form 20-F for the year ended December 31, 2024, filed with the U.S. Securities and Exchange Commission, the events and circumstances discussed in such forward-looking statements may not occur, and Pharming's actual results could differ materially and adversely from those anticipated or implied thereby. All forward-looking statements contained in this presentation are expressly qualified in their entirety by the cautionary statements contained or referred to in this section. Readers should not place undue reliance on forward-looking statements. Any forward-looking statements speak only as of the date of this presentation and are based on information available to Pharming as of the date of this presentation. Pharming does not undertake any obligation to publicly update or revise any forward-looking statement as a result of new information, future events or other information.*

# History of growth and innovation at Pharming



***Develop a leading global rare disease company  
with a diverse portfolio and presence in large markets,  
leveraging proven and efficient clinical development,  
supply chain, and commercial infrastructure***



## **Revenues**

**FY24: US\$297 million (+21%)**

**4Q24: US\$93 million (+14%)**

**Operating profit and  
positive operating cash flow in 3Q & 4Q 2024**

## RUCONEST®

### ◆ Revenue:

FY24 US\$252.2M (+11%)

4Q24 US\$79.6M (+9%)

### ◆ Strong U.S. in-market demand

U.S. physician prescriber base +11% FY24

New enrollments up 24% FY24

## Joenja®

### ◆ Revenue:

FY24 US\$45.0M (+147%)


4Q24 US\$13.1M (+65%)

### ◆ Increasing APDS patients on therapy

Found >240 in the U.S. and >880 globally

Paid therapy: 96 patients + 5 pending (U.S.)

Additional 188 patients on therapy globally  
(access programs and clinical studies)







**RUCONEST® for HAE**




# RUCONEST® (rhC1INH):

## 2<sup>nd</sup> most prescribed therapy for acute HAE attacks in the US

Only recombinant treatment that targets the root cause of HAE by replacing C1-INH

Only recombinant treatment that acts at multiple points in the cascades leading to HAE attacks




◆ Type 1, Type 2, and Normal C1-INH HAE patients rely on RUCONEST

◆ 97% patients needed just 1 dose<sup>1</sup>


◆ 93% acute attacks stopped for at least 3 days<sup>2</sup>

◆ RUCONEST mostly used by patients experiencing moderate to severe attacks, who attack more frequently


- Fail on icatibant and other acute therapies
- Need to re-dose with other treatments to resolve attacks



Time of taking  
RUCONEST



4 hours after



24 hours after






Pharming<sup>®</sup>

**Joenja<sup>®</sup> (leniolisib) for APDS**  
**leniolisib for PIDs with Immune**  
**Dysregulation**

# APDS is a rare primary immunodeficiency (PID)

## Genetic defect leads to PI3K $\delta$ hyperactivity


Hyperactive PI3K $\delta$  results in dysregulated B and T cell development<sup>1-3</sup>



The PI3K $\delta$  enzyme is at the beginning of a complex signaling pathway

Note: Illustration does not include all steps in the signaling pathway.

Immune imbalance leads to diverse signs and symptoms<sup>1,4-6</sup>



### Severe, recurrent, persistent infections

- Sinopulmonary
- Herpesvirus (especially EBV and CMV)

### Lymphoproliferation

- Lymphadenopathy
- Splenomegaly/hepatomegaly
- Nodular lymphoid hyperplasia

### Enteropathy

### Autoimmunity

- Cytopenias
- Autoimmune disorders
- Autoinflammatory disorders

### Bronchiectasis

### Lymphoma

FOXO, forkhead box O; mTOR, mammalian target of rapamycin; PI3K $\delta$ , phosphoinositide 3-kinase delta; PKB, protein kinase B.

1. Lucas CL, et al. *Nat Immunol*. 2014;15(1):88-97. 2. Fruman DA, et al. *Cell*. 2017;170(4):605-635. 3. Okkenhaug K, Vanhaesebroeck B. *Nat Rev Immunol*. 2003;3(4):317-330. 4. Coulter TI, et al. *J Allergy Clin Immunol*. 2017;139(2):597-606. 5. Elkaim E, et al. *J Allergy Clin Immunol*. 2016;138(1):210-218. 6. Jamee M, et al. *Clin Rev Allergy Immunol*. 2020;59(3):323-333.



# Joenja<sup>®</sup>: First and only approved therapy for APDS



Joenja<sup>®</sup> (leniolisib) is an oral medication used to treat activated phosphoinositide 3-kinase delta (PI3K $\delta$ ) syndrome (APDS) in adult and pediatric patients 12 years of age and older

Joenja<sup>®</sup> targets the root cause of APDS

- Normalizes the hyperactive PI3K $\delta$  pathway to correct the underlying immune defect in APDS patients
- Helps address both immune deficiency and immune dysregulation




No drug-related serious adverse events or study withdrawals in Joenja<sup>®</sup> trials  
Clinical data and tolerability for long term treatment

Approved in the US (Mar 2023), Israel (Apr 2024), UK (Sept 2024), Australia (Mar 2025)

Regulatory reviews on-going in the EU, Canada and several other countries

Submission planned in Japan in 2025

# Joenja<sup>®</sup> (leniolisib) – Reaching more APDS patients and expanding the addressable patient population



\* 96 patients on paid therapy + 5 pending. U.S. Pricing: 30-day supply \$49,500, Annual cost (WAC) \$594,000



## Variants of Uncertain Significance

- ❖ VUSs: insufficient data to determine if variant is disease causing
- ❖ >1200 patients in the U.S.
- ❖ VUSs may be reclassified as APDS with additional evidence\*



## VUS study results

- ❖ High throughput screening (MAVE) study, completed in December, identified novel variants leading to PI3K $\delta$  hyperactivity
- ❖ Genetics testing labs to review study data, reclassify variants and update test reports
- ❖ Additional APDS patients to be identified over the course of 2025

\* As results become available, patients with validated variants could be diagnosed with APDS and be eligible for Joenja® treatment.




## Pediatric

Phase III trial for children 4-11 years old with APDS

### **Positive topline data announced December 2024**

- ◆ 21 patients enrolled in U.S., Europe, and Japan
- ◆ Both co-primary endpoints show improvement consistent with the RCT in adolescents and adults
- ◆ Benefits seen across the four tested dose levels
- ◆ No deaths/discontinuations due to AEs. No new safety findings
- ◆ Data to be presented at CIS conference in May
- ◆ Regulatory filings beginning with the U.S. in second half 2025

# Joenja<sup>®</sup> (leniolisib) – Reaching more APDS patients and expanding the addressable patient population



\* 96 patients on paid therapy + 5 pending. U.S. Pricing: 30-day supply \$49,500, Annual cost (WAC) \$594,000

## Two Phase II studies underway to target PI3Kδ


- Critical role of PI3Kδ in lymphocyte regulation
- Patient manifestations with similarities to APDS and large unmet clinical need
- Therapeutic strategy: modulate PI3Kδ to address lymphoproliferation and autoimmunity

## Genetically defined PIDs with immune dysregulation linked to PI3Kδ signaling<sup>1</sup>

- Phase II study started Oct 2024<sup>2</sup>
- N=12 patients, treated for 20 weeks
- FDA Fast Track designation
- Conducted at NIH

## Common variable immunodeficiency (CVID) with immune dysregulation

- Phase II study started Feb 2025<sup>3</sup>
- N=20 patients, treated for 24 weeks




*Not to scale with population sizes*

1. PIDs include ALPS-FAS, CTLA4 haploinsufficiency, NFKB1 haploinsufficiency and PTEN deficiency, amongst others

2. Single arm, open-label, dose range-finding study. ClinicalTrials.gov ID NCT06549114

3. Single arm, open-label, dose range-finding study. ClinicalTrials.gov ID NCT06897358

## Phase II proof of concept clinical trial – single arm, open-label, dose range-finding study (N=20)



- Multi-center study (US, UK, EU)
- Patients with a CVID diagnosis, evidence of lymphoproliferation, and at least one additional clinical manifestation of immune dysregulation
- Primary: Safety & Tolerability
- Secondary/Exploratory: PK/PD, efficacy measures
- 10/30/70 mg BID: 4/4/16 wks treatment, resp.
- Inform dose regimen/design of Phase III program

Beth Israel Lahey Health   
**Lahey Hospital & Medical Center**

Lead Investigator:

Jocelyn Farmer, MD, PhD  
Director of the Clinical Immunodeficiency  
Program



Pharming®

**KL1333 for mtDNA**  
**Mitochondrial Disease**



## Primary mitochondrial diseases – rare disorders impairing mitochondrial energy production

- Severe fatigue, myopathy, and reduced life expectancy
- Poor quality of life (e.g., loss of job, social isolation, depression)



## KL1333 positioned to become first standard of care in mitochondrial DNA disease

- Novel mechanism of action addresses the underlying disorder
- >30,000 diagnosed patients\*



## Pivotal study ongoing with positive interim analysis

- Patient recruitment for second wave of pivotal FALCON clinical trial to start shortly
- Read-out anticipated in 2027 with potential FDA approval by end of 2028



## Significant unmet medical need and no approved therapies

- Builds on Pharming's existing rare disease expertise and infrastructure
- Concentrated centers of excellence and strong advocacy groups

\* in US, EU4 and UK

## Pivotal FALCON Study

### WAVE 1 – Fully enrolled

- ◆ 40 patients recruited across six countries (U.S., UK, France, Spain, Belgium, Denmark)
- ◆ 18 sites activated
- ◆ Interim analysis at 24 weeks conducted in Q3 2024

### WAVE 2 – Expansion

- ◆ 180 total patients treated for 48 weeks
  - Wave 1 sites ready to start enrolling
  - Wave 2 sites undergoing activation
- ◆ Readout anticipated 2027

### Interim Futility Analysis:

*Positive outcome achieved, with both primary endpoints having passed futility*

- ◆ Promising differences favoring the active arm vs. placebo for both primary efficacy endpoints; if trends continue consistently, we expect a successful result at the completion of this trial
- ◆ Data monitoring committee (DMC) recommended continuing with Wave 2:
  - Safety and tolerability profile acceptable
  - No changes to study design
  - 180 total patients confirmed in the study





## Growing commercial portfolio

- RUCONEST's specific positioning within the on-demand HAE market
- Joenja (leniolisib) for APDS: VUSs, pediatric label, geographic expansion



## High value pipeline

- Leniolisib new indications (PIDs with immune dysregulation)
- Abliva KL1333 (mtDNA mitochondrial disease)



## Organizational efficiency

- Targeted geographic expansion (8 key markets) with WW access program
- Efficient and scalable organization for portfolio development



**Revenues:**  
**2024: US\$297 million (21% growth)**  
**2025 guidance: US\$315 - 335 million**

**EURONEXT AMS: PHARM**  
**Nasdaq: PHAR**



[www.pharming.com](http://www.pharming.com)

NASDAQ: **PHAR** | EURONEXT Amsterdam: **PHARM**




**Pharming Group N.V.**

# Appendix

# APDS can impact many facets of life


## Physical<sup>1,2</sup>

Frequent infections  
Swollen glands  
Shortness of breath  
Coughing/wheezing  
Chest or joint pain  
Fatigue  
Inability to exercise  
Hearing loss  
Diarrhea  
Skin problems




## Social<sup>3,4</sup>

Missing school, work, or daily activities



## Treatment Burden<sup>1-4</sup>

Frequent hospitalizations  
Surgeries  
Visiting multiple doctors  
Invasive or time-consuming treatments




## Mental<sup>1,3-5</sup>

Anxiety  
Depression  
Stress




# Heterogeneous, evolving symptomology can often lead to missed diagnoses

## Timeline of the most common pathologies\* seen in APDS<sup>1-4</sup>



APDS has often been diagnosed as another PI or condition, causing delays in diagnosis<sup>1</sup>



Improved identification of symptoms, increased genetic testing, and earlier diagnosis are needed

\*Pathologies can occur at any time.

<sup>†</sup>In Elkaim APDS2 cohort, median age of bronchiectasis is 13; in Maccari ESID cohort, median age is 11.2.


<sup>‡</sup>No median ages are available for these manifestations.

ALPS, autoimmune lymphoproliferative syndrome; CID, combined immunodeficiency; CVID, common variable immune deficiency; ESID, European Society for Immunodeficiencies; HIGM, hyper immunoglobulin M syndrome; IgG, immunoglobulin G; PI3Kδ, phosphoinositide 3-kinase delta; XLA, X-linked agammaglobulinemia.

1. Jamee M, et al. *Clin Rev Allergy Immunol*. 2020;59(3):323-333. 2. Maccari ME, et al. *Front Immunol*. 2018;9:543. 3. Elkaim E, et al. *J Allergy Clin Immunol*. 2016;138(1):210-218.e9. 4. Coulter TI, et al. *J Allergy Clin Immunol*. 2017;139(2):597-606.

## Immune Deficiency

- Antimicrobial prophylaxis
- Immunoglobulin replacement therapy



## Immune Dysregulation

- Corticosteroids
- Other immunosuppressants
- mTOR inhibitors

*None of these therapies are FDA-approved for APDS treatment*

Hematopoietic stem cell transplant

APDS, activated phosphatidylinositol 3-kinase  $\delta$  syndrome; IRT, immunoglobulin replacement therapy; mTOR, mammalian target of rapamycin; PI, primary immunodeficiency; PIRD, primary immune regulatory disorder.

1. Coulter TI, et al. *J Allergy Clin Immunol.* 2017;139(2):597-606. 2. Elkaim E, et al. *J Allergy Clin Immunol.* 2016;138(1):210-218. 3. Chan AY, et al. *Front Immunol.* 2020;11:239. 4. Chinn IK, et al. *J Allergy Clin Immunol.* 2020;145(1):46-69.

## Pivotal Trial - Part 1: Dose- finding<sup>1,2</sup>



Nonrandomized, open-label,  
dose-escalating



6 patients with APDS



12 weeks



10 mg, 30 mg, 70 mg bid  
(4 weeks each dose)



70 mg bid selected for Part 2

## Pivotal Trial - Part 2: Efficacy & Safety Evaluation<sup>3</sup>



Randomized, triple-blinded,  
placebo-controlled



31 patients with APDS  
(21 Joenja®, 10 placebo)



12 weeks



70 mg bid



Co-primary efficacy end points

- Change from baseline in log<sup>10</sup>-transformed SPD of index lesions
  - Also assessed as % change
- Change from baseline in percentage of naïve B cells out of total B cells

Secondary and exploratory end points

Safety

## Open-label extension study<sup>4,5</sup>



Nonrandomized, open-label,  
long-term study



• 35 patients with APDS from  
Parts 1 and 2

• 2 patients with APDS previously  
treated with investigational  
PI3Kδ inhibitors



Ongoing



70 mg bid



Long-term safety, tolerability,  
efficacy, and pharmacokinetics

bid, twice a day; PI3Kδ, phosphoinositide 3-kinase delta; SPD, sum of product diameters

1. Rao VK, et al. *Blood*. 2017;130(21):2307-2316. 2. NCT02435173. ClinicalTrials.gov. <https://clinicaltrials.gov/ct2/show/NCT02435173>. Updated May 6, 2015. Accessed March 13, 2023. 3. Rao VK, et al. *Blood*. 2023;141(9):971-983.


4. NCT02859727. ClinicalTrials.gov. <https://clinicaltrials.gov/ct2/show/NCT02859727>. Updated October 31, 2022. Accessed March 3, 2023. 5. Data on file. Pharming Healthcare Inc; 2022.




# Joenia® addresses the underlying cause of APDS to help restore immune balance – Phase 3 co-primary endpoints

## At 12 weeks Joenia® decreased lymphadenopathy and increased naïve B cells

### Immune Dysregulation



### Immune Deficiency



Data were analyzed using an ANCOVA model with treatment as a fixed effect and baseline as a covariate. Use of glucocorticoids and IRT at baseline were both included as categorical (Yes/No) covariates. Baseline is defined as the arithmetic mean of the baseline and D1 values when both are available, and if either baseline or the D1 value is missing, the existing value is used. P-value is 2-sided. Least square means are graphed. Error bars are standard error of the mean.

\*The analysis excluded 2 patients from each treatment group due to protocol deviations and 1 Joenia patient having complete resolution of the index lesion identified at baseline.

†Out of 27 patients in the PD analysis set, 13 patients met the analysis requirements, including having a percentage of <48% of naïve B cells at baseline, to form the B-PD analysis set.

Joenia [package insert]. Leiden, The Netherlands: Pharming Technologies B.V.; 2023.

## Secondary endpoint: Significant reductions in spleen size by 2D and 3D analysis compared to placebo

- The adjusted mean difference in bidimensional spleen size between Joenia<sup>®</sup> (n=19) and placebo (n=9) was  $-13.5 \text{ cm}^2$  (95% CI:  $-24.1, -2.91$ ),  $P=0.0148$
- The adjusted mean difference in 3D spleen volume between Joenia<sup>®</sup> (n=19) and placebo (n=9) was  $-186 \text{ cm}^3$  (95% CI:  $-297, -76.2$ ),  $P=0.0020$

at week 12

**27%**

reduction in 3D spleen volume\*

Secondary measure: spleen volume scan results of actual patient illustrate average improvement documented for patients taking Joenia<sup>®</sup>

Prior to treatment:  
491 mL



At week 12:  
314 mL



Actual patient images of a 17-year-old male. As individual results vary, images may not be representative of all patients.


Rao VK, et al. Blood. 2023;141(9):971-983.

\*In the PD analysis set, the mean (SD) percentage change from baseline to week 12 in 3D spleen volume ( $\text{mm}^3$ ) was  $-26.68\%$  (12.137) with Joenia<sup>®</sup> (n=19) and  $-1.37\%$  (24.238) with placebo (n=9). The ANCOVA model was used with treatment as a fixed effect and  $\log_{10}$ -transformed baseline as a covariate for index and non-index lesions. The use of both glucocorticoids and IV Ig at baseline was included as categorical (yes/no) covariates.

This analysis excluded 2 patients in each treatment group. In the Joenia<sup>®</sup> group, 1 patient with a complete index lesion response was excluded, and 3 patients were excluded for no non-index lesion at baseline. PD, pharmacodynamics.

# An exploratory endpoint showed Joenja® reduced IgM levels

## Mean serum IgM rapidly reduced to within normal limits



- In the Joenja® arm, IgM was elevated above normal limits in 6 patients at baseline, and by week 12 was reduced in all, with 50% returning to within normal limits
- In contrast, IgM was elevated above normal limits at baseline in 4 patients in the placebo arm, and by week 12 levels remained stable or elevated, with 0% returning to within normal limits

Error bars are standard error of the mean. Safety analysis set (N=31) shown. Blue box indicates IgM normal range.

Soluble biomarkers, including IgM, were prespecified exploratory endpoints in the protocol. Although an observational decrease in IgM was noted in some patients, no statistical significance can be made from this analysis, and no conclusions should be drawn.

Rao VK, et al. Blood. 2023;141(9):971-983

## Phase 3 Trial<sup>1,2</sup>

Adverse reactions reported by ≥2 patients treated with Joenja and more frequently than placebo

	Joenja (n=21) n (%)	Placebo (n=10) n (%)
Headache	5 (24)	2 (20)
Sinusitis	4 (19)	0
Dermatitis atopic*	3 (14)	0
Tachycardia <sup>†</sup>	2 (10)	0
Diarrhea	2 (10)	0
Fatigue	2 (10)	1 (10)
Pyrexia	2 (10)	0
Back pain	2 (10)	0
Neck pain	2 (10)	0
Alopecia	2 (10)	0

- Study drug-related AEs occurred in 8 patients; the incidence was lower in the Joenja arm (23.8%) than in the placebo arm (30.0%)
- No AEs led to discontinuation of study treatment

A patient with multiple occurrences of an AE is counted only once in the AE category. Only AEs occurring at or after first drug intake are included.

\*Includes dermatitis atopic and eczema. <sup>†</sup>Includes tachycardia and sinus tachycardia.

AEs, adverse events; ALT, alanine aminotransferase; AST, aspartate aminotransferase; SAE, serious adverse event.

1. Rao VK, et al. Blood. 2023;141(9):971-983. 2. Joenja [package insert]. Leiden, The Netherlands: Pharming Technologies B.V.; 2023. 3. Data on file. Pharming Healthcare Inc; 2022.

Please see Important Safety Information and full Prescribing Information available at joenja.com

## Open-label Extension Study<sup>3</sup>

Data cutoff for interim analysis: December 13, 2021



- 32/37 patients reported ≥1 AE
- 78.4% of AEs were grade 1, 48.6% grade 2, 27.0% grade 3, 0% grade 4
- No SAEs related to Joenja

Most common AEs	n
Upper respiratory tract infection	8
Headache	6
Pyrexia	6
Otitis externa	5
Weight increase	5
COVID-19, positive/negative	5/14

One patient with significant baseline cardiovascular comorbidities suffered cardiac arrest resulting in death at extension Day 879; determined by investigator not to be related to study drug

- Across all trials<sup>2</sup>**
- 38 patients had a **median exposure of ~2 years**
  - 4 patients had **>5 years of exposure**

# Open-label extension interim analysis of days spent with infections and IRT reduction



Although safety was the primary objective of the open-label study, this post hoc analysis from the open-label study was not powered to provide any statistical significance of efficacy and therefore no conclusions should be drawn.

\*Infections that developed during the study were reported as adverse events. Investigators were requested to inquire about signs and symptoms of infections at each visit, with a particular focus on bacterial enterocolitis. Patients were not provided an infection diary to document infections occurring between visits. One patient was excluded from the analysis due to an incorrect year that was recorded for an infection.

<sup>†</sup>Baseline infections are each group's year 1 annual rate of infections. N values changed because patients were in the OLE for different lengths of time. <sup>‡</sup>Data on concomitant medication usage was reported at each patient visit.

<sup>§</sup>One patient had a subsequent one-time dose. <sup>||</sup>One patient achieved IRT freedom for 3 months but subsequently restarted IRT. IRT, immunoglobulin replacement therapy; m, number of infection days; N, number of patients in follow-up category.

Rao VK, et al. Poster presented at: 64<sup>th</sup> Annual American Society of Hematology Annual Meeting; December 10-13, 2022; New Orleans, LA.

Please see Important Safety Information and full Prescribing Information available at [joenja.com](http://joenja.com)

## VUSs frustrate patients and doctors, limiting diagnosis of genetic diseases such as APDS



Pharming is aware of **~1,200 US patients** harboring *PIK3CD/R1* VUSs

- This figure will continue to grow over time
- VUS are identified at ~4x the rate of likely pathogenic/pathogenic (LP/P) variants
- Similar VUS frequencies expected worldwide
- Published literature, which includes more than 1.5 million patients, showed that 20% of reclassified VUSs are upgraded to LP/P
- Pilot study in 25 VUS patient samples - findings consistent with APDS identified in 5 patients (20%) including patient preparing for enrollment


**No systemic initiatives exist to resolve *PIK3CD/R1* VUSs, yet these patients remain a significant opportunity to identify incremental patients with APDS**




# Given importance of PI3K $\delta$ in B & T cells, immune dysregulation in PIDs can occur via alterations in PI3K $\delta$ signaling

## Altered PI3K $\delta$ signaling can occur in multiple PID genetic disorders beyond APDS

(ex: ALPS-FAS, CTLA4, NFKB1, PTEN)<sup>1-4</sup>



### leniolisib




**High unmet medical need**  
- no approved therapies other than Joenja<sup>®</sup> (leniolisib) for APDS:  
SOC immunosuppressives (e.g. rapamycin) have limited efficacy and significant tolerability concerns

## Clinical manifestations, disease onset and severity similar to APDS<sup>5-10</sup>




### Lymphoproliferation

- Lymphadenopathy
- Splenomegaly/hepatomegaly
- Nodular lymphoid hyperplasia



### Autoimmunity

- Cytopenias
- Autoimmune disorders
- Autoinflammation



### GI Disease

- Autoimmune enteropathy
- Nodular regenerative hyperplasia



### Pulmonary Disease

- GLILD
- Bronchiectasis



### Infections

- Sinopulmonary
- Herpesvirus



### Lymphoma

Note: Illustration does not include all steps in the signaling pathway.

FOXO, forkhead box O; mTOR, mammalian target of rapamycin; PI3K $\delta$ , phosphoinositide 3-kinase delta; PKB, protein kinase B.


1. Volkl et al. Blood 2016; 128(2):227-238. 2. Tsujita, et al. J Allergy Clin Immunol. 2016;138(6):1872-80. 3. Rowshanravan B, et al. Blood. 2018;131(1):58-67. 4. Additional unpublished collaborator data. 5. Bride K & Teachey D. F1000Res. 2017;6:1928

6. Kuehn HS, et al. Science 2014; 345:1623-27. 7. Lorenzini T, et al. J Allergy Clin Immunol. 2020;146:901-11. 8. Eissing, et al. Transl Oncol. 2019;12(2):361-3672. 9. Coulter TI, et al. J Allergy Clin Immunol. 2017;139(2):597-606. 10. Schwab C, et al. J

Allergy Clin Immunol. 2018;142(6):1932-1946.



## Phase II proof of concept clinical trial – single arm, open-label, dose range-finding study (N=12)



- Patients with PIDs linked to PI3K $\delta$  signaling, e.g. ALPS-FAS<sup>1</sup>, CTLA4 haploinsufficiency<sup>2</sup>, NFKB1 haploinsufficiency<sup>3</sup>, PTEN deficiency<sup>4</sup> (treatable population ~7.5/million)
- Primary: Safety & Tolerability
- Secondary/Exploratory: PK/PD, efficacy measures
- 10/30/70 mg: 4/4/12 wks treatment, respectively
- Pick Best Dose regimen for Phase III



National Institute of  
Allergy and  
Infectious Diseases

Lead Investigator: Gulbu Uzel, M.D., Senior  
Research Physician


Co-Investigator: V. Koneti Rao, M.D., FRCPA,  
Senior Research Physician  
Primary Immune Deficiency Clinic (ALPS  
Clinic)

1. Bride K & Teachey D. F1000Res. 2017;6:1928. ; Rao VK & Oliveria JB. Blood 2011; 118(22):5741-51.  
2. Kuehn HS, et al. Science 2014; 345:1623-27. ; Schwab C, et al. J Allergy Clin Immunol. 2018;142(6):1932-1946.  
3. Lorenzini T, et al. J Allergy Clin Immunol. 2020;146:901-11.  
4. Eissing M, et al. Transl Oncol. 2019;12(2):361-367. ; Tsujita, et al. J Allergy Clin Immunol. 2016;138(6):1872-80.

## Primary Mitochondrial Disease (PMD)

- ❖ Mitochondria, often described as the “powerhouses” of cells, are crucial for energy production
- ❖ Mitochondrial diseases are a group of genetic disorders characterized by dysfunctional mitochondria due to mutations in mitochondrial (mtDNA) or nuclear DNA
- ❖ The abnormal  $\text{NAD}^+/\text{NADH}$  ratio results in decreased ATP production, contributing to organ dysfunction and disease deterioration
- ❖ For patients this means symptoms of severe fatigue and muscle weakness – symptoms which patients report as the most troublesome\*

### Dysfunctional Mitochondria



- ↓ Decreased energy production
- ↓ Decreased mitochondria biogenesis

\*Voice of the Patient Report, United Mitochondrial Disease Foundation, 2019.

## Presentation and Diagnosis

- ◆ Patients present to their primary care doctor and then often get referred to a neurologist for musculoskeletal issues
- ◆ Either the neurologist or a referral to a metabolic geneticist will result in a diagnosis
- ◆ Many patients are diagnosed at academic centers specializing in mitochondrial disease
- ◆ A combination of routine lab tests and genetic testing available from major testing labs help to diagnose patients

## Impact

- ◆ Patients heavily burdened in their daily lives including symptoms like severe fatigue, myopathy, and metabolic dysfunction
- ◆ Impact on QoL including loss of job, loss of independence, depression/anxiety
- ◆ Primary mitochondrial diseases lead to a three-to-four-decade reduction in life-expectancy

## Treatment


- ◆ No approved treatment options
- ◆ Patients are limited to using vitamins, supplements, and physical therapy

***“On the worst days I will be crying in frustration because going to the kitchen seems equivalent to climbing a mountain and just trying to process what others are saying to me involves all the energy and concentration that I have.”***


United Mitochondrial Disease Foundation, Voice of the Patient Conference, 2019

## KL1333 normalizes conversation of NADH to NAD<sup>+</sup> via NQO1

### Normalizes the NAD<sup>+</sup>/NADH Ratio



### Restored Energy Metabolism



## Attributes

- ◆ Directly increases the NAD<sup>+</sup>/NADH ratio via NQO1
- ◆ Unique MoA works upstream from all competing MoA in PMD
- ◆ Oral, small molecule, BID dosing
- ◆ Favourable safety profile
- ◆ Favourable IP protection
- ◆ Orphan Drug Designation in US & EU and FDA Fast Track
- ◆ Potential first-in-disease with registrational clinical study




## Outcomes

- ◆ Improved energy regulation and ETC function
- ◆ Stimulation of mitochondria biogenesis
- ◆ Fatigue reduction
- ◆ Increased exercise capacity

# Phase 1b demonstrated significant activity vs. placebo

## The placebo-controlled Phase 1b study demonstrated that KL1333 reduced patients' fatigue and myopathy after only 10 days, 50 mg/day

- ◆ KL1333 demonstrated efficacy in the phase 1b placebo-controlled portion with patients diagnosed with mtDNA mitochondrial disease
  - Fatigue reduction (NeuroQoL fatigue change)
  - Muscle function improvement (30 seconds sit-to-stand)
- ◆ KL1333 showed efficacy signals after 10 days using 50 mg/day
- ◆ Mitochondrial patients have increased lactate levels and increasing the concentration of KL1333 resulted in an improved lactate/pyruvate ratio, reflecting target engagement
- ◆ No serious adverse events reported



## Regulatory Feedback

- ◆ Both FDA and EMA accepted study as registrational
- ◆ FDA said achieving one of the two endpoints would be sufficient for filing
- ◆ Conducted regular and detailed discussions with the FDA to facilitate alignment

## Study Design


- ◆ **Methodology**
  - Randomized, double-blind, parallel-group, placebo-controlled pivotal study
- ◆ **Patients Included**
  - Adult PMD patients with mtDNA mutations\* with fatigue and myopathy
- ◆ **Primary Endpoints**
  - Fatigue using the PROMIS Fatigue Mitochondrial Disease Short Form
  - Muscle weakness using the 30 second Sit-to-Stand test

**Week 24**  
Interim futility analysis  
(Wave 1 only)

**Week 48**  
Primary efficacy analysis

**Week 53**  
Safety follow-up


## Study Schematic



\*Most prevalent mtDNA disorders include m.3243A>G associated MELAS-MIDD spectrum disorders, single large scale mtDNA deletion associated KSS-CPEO spectrum disorders, other multisystemic mtDNA-related disease (including MERRF)



# Significant revenue opportunity for KL1333




***>30,000 diagnosed mtDNA mitochondrial disease patients addressable in the US, EU4 and UK***

\*mtDNA mutations including m.8344A>G MELAS-MIDD, MERRF, KSS-CEPO, large scale mtDNA deletions

<sup>1</sup>Gorman, G.S. et al. Prevalence of nuclear and mitochondrial DNA mutations related to adult mitochondrial disease. Ann Neurol 2015 May;77(5):753-9.







<sup>2</sup>Gorman, G.S. et al. Mitochondrial Diseases. Nat. Rev. Vol 2, 1-22 (2016).

# Majority of patients diagnosed and treated in US Centers of Excellence or academic institutions



# Other programs focus on different patient population or failed with different MOA

*Previous programs failed due to old mechanisms of action or evaluating the wrong endpoints*

Asset	Type	MOA / ROA	Stage	Patient Group	Comments
 <b>KL1333</b>	Small molecule	NAD <sup>+</sup> /NADH modulator Oral	Pivotal	mtDNA mutations (e.g., mtDNA deletion, m.8344A>G, MELAS-MIDD, MERRF, KSS-CEPO)	<ul style="list-style-type: none"> <li>▪ Ongoing potentially registrational phase 2 study</li> <li>▪ FALCON pivotal study reported positive 24w interim analysis</li> </ul>
 <b>Elamipretide</b>	Peptide	Cardiolipin stabilizer Subcutaneous	Phase 3	nDNA mutations	<ul style="list-style-type: none"> <li>▪ nDNA represents about 20% of PMD patients</li> <li>▪ In discussions with FDA for ultra rare Barth syndrome</li> </ul>
 <b>Zagociguat</b>	Small molecule	Guanylate cyclase stimulator Oral	Phase 2b ready	MELAS	<ul style="list-style-type: none"> <li>▪ Completed open-label MELAS phase 2a</li> <li>▪ Phase 2b trial planned with focus on fatigue, myopathy and cognition</li> </ul>
 <b>Sonlicromanol</b>	Small molecule	Redox modulator Oral	Phase 3 ready	mtDNA mutation (MELAS-MIDD)	<ul style="list-style-type: none"> <li>▪ Phase 2a study in m.3243A&gt;G patients showed predominantly neutral results across multiple endpoints</li> <li>▪ Phase 2b study failed primary endpoint, positive changes in post-hoc analyses and open-label extension</li> </ul>
 <b>Mavodelpar</b>	Small molecule	PPAR $\delta$ agonist Oral	NA	mtDNA in the interventional trial and extended to include nDNA in the OLE	Phase 3 failed to achieve primary endpoint of 12-minute walk test
 <b>Boicedelpar</b>	Small molecule	PPAR $\delta$ agonist Oral	NA	Mixed population of mtDNA and nDNA	Phase 2 program using 6-minute walk test terminated